

## Product datasheet for RC210469L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MCP3 (CCL7) (NM\_006273) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** MCP3 (CCL7) (NM\_006273) Human Tagged ORF Clone Lentiviral Particle

Symbol: MCP3

Synonyms: FIC; MARC; MCP-3; MCP3; NC28; SCYA6; SCYA7

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 006273

ORF Size: 297 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC210469).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 006273.2

 RefSeq Size:
 815 bp

 RefSeq ORF:
 300 bp

 Locus ID:
 6354

 UniProt ID:
 P80098

 Cytogenetics:
 17q12

**Protein Families:** Druggable Genome, Secreted Protein





## MCP3 (CCL7) (NM\_006273) Human Tagged ORF Clone Lentiviral Particle - RC210469L3V

**Protein Pathways:** Chemokine signaling pathway, Cytokine-cytokine receptor interaction, NOD-like receptor

signaling pathway

MW: 11.2 kDa

**Gene Summary:** This gene encodes monocyte chemotactic protein 3, a secreted chemokine which attracts

macrophages during inflammation and metastasis. It is a member of the C-C subfamily of chemokines which are characterized by having two adjacent cysteine residues. The protein is an in vivo substrate of matrix metalloproteinase 2, an enzyme which degrades components of the extracellular matrix. This gene is part of a cluster of C-C chemokine family members on

chromosome 17q. [provided by RefSeq, Jul 2008]