

Product datasheet for RC210483L2

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MC2 receptor (MC2R) (NM_000529) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: MC2 receptor (MC2R) (NM_000529) Human Tagged Lenti ORF Clone

Tag: mGFP

Symbol: MC2 receptor

Synonyms: ACTHR
Mammalian Cell None

Selection:

Vector: pLenti-C-mGFP (PS100071)

E. coli Selection: Chloramphenicol (34 ug/mL)


ORF Nucleotide The ORF insert of this clone is exactly the same as(RC210483).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_000529

ORF Size: 891 bp





OTI Disclaimer:

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at customercom or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

OTI Annotation:

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:

Cytogenetics:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: NM 000529.2, NP 000520.1

 RefSeq Size:
 3652 bp

 RefSeq ORF:
 894 bp

 Locus ID:
 4158

 UniProt ID:
 Q01718

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Neuroactive ligand-receptor interaction

18p11.21

MW: 33.9 kDa


Gene Summary: MC2R encodes one member of the five-member G-protein associated melanocortin receptor

family. Melanocortins (melanocyte-stimulating hormones and adrenocorticotropic hormone) are peptides derived from pro-opiomelanocortin (POMC). MC2R is selectively activated by adrenocorticotropic hormone, whereas the other four melanocortin receptors recognize a variety of melanocortin ligands. Mutations in MC2R can result in familial glucocorticoid deficiency. Alternate transcript variants have been found for this gene. [provided by RefSeq,

May 2014]



Product images:



Circular map for RC210483L2