

Product datasheet for **RC210533L2V**

ER81 (ETV1) (NM_004956) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	ER81 (ETV1) (NM_004956) Human Tagged ORF Clone Lentiviral Particle
Symbol:	ER81
Synonyms:	ER81
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_004956
ORF Size:	1431 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC210533).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_004956.3
RefSeq Size:	6824 bp
RefSeq ORF:	1434 bp
Locus ID:	2115
UniProt ID:	P50549
Cytogenetics:	7p21.2
Domains:	ETS, ETS_PEA3_N
Protein Families:	ES Cell Differentiation/IPS, Transcription Factors



[View online »](#)

MW: 55.1 kDa

Gene Summary: This gene encodes a member of the ETS (E twenty-six) family of transcription factors. The ETS proteins regulate many target genes that modulate biological processes like cell growth, angiogenesis, migration, proliferation and differentiation. All ETS proteins contain an ETS DNA-binding domain that binds to DNA sequences containing the consensus 5'-CGGA[AT]-3'. The protein encoded by this gene contains a conserved short acidic transactivation domain (TAD) in the N-terminal region, in addition to the ETS DNA-binding domain in the C-terminal region. This gene is involved in chromosomal translocations, which result in multiple fusion proteins including EWS-ETV1 in Ewing sarcoma and at least 10 ETV1 partners (see PMID: 19657377, Table 1) in prostate cancer. In addition to chromosomal rearrangement, this gene is overexpressed in prostate cancer, melanoma and gastrointestinal stromal tumor. Multiple alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2016]