

## Product datasheet for RC210546L1

### VCIP 135 (VCPIP1) (NM\_025054) Human Tagged Lenti ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	VCIP 135 (VCPIP1) (NM_025054) Human Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	VCIP 135
Synonyms:	DUBA3; VCIP135
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC210546).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.


ACCN:	NM_025054
ORF Size:	3666 bp




[View online >](#)

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_025054.3</a>
<b>RefSeq Size:</b>	8192 bp
<b>RefSeq ORF:</b>	3669 bp
<b>Locus ID:</b>	80124
<b>UniProt ID:</b>	<a href="#">Q96JH7</a>
<b>Cytogenetics:</b>	8q13.1
<b>MW:</b>	134.1 kDa
<b>Gene Summary:</b>	Acts as a deubiquitinating enzyme. Necessary for VCP-mediated reassembly of Golgi stacks after mitosis. May play a role in VCP-mediated formation of transitional endoplasmic reticulum (tER). Mediates dissociation of the ternary complex containing STX5A, NSFL1C and VCP (By similarity). Hydrolyzes 'Lys-11'- and 'Lys-48'-linked polyubiquitin chains. [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RC210546L1



Double digestion of RC210546L1 using SgfI and MluI