

Product datasheet for RC210559L3

SPDYA (NM_182756) Human Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	SPDYA (NM_182756) Human Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	SPDYA
Synonyms:	RINGO3; RINGOA; SPDY1; SPY1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC210559).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_182756
ORF Size:	939 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_182756.2
RefSeq Size:	1830 bp
RefSeq ORF:	942 bp
Locus ID:	245711
UniProt ID:	Q5MJ70
Cytogenetics:	2p23.2
Protein Pathways:	Oocyte meiosis, Progesterone-mediated oocyte maturation
MW:	36.5 kDa
Gene Summary:	Regulates the G1/S phase transition of the cell cycle by binding and activating CDK1 and CDK2 (PubMed:12972555). Contributes to CDK2 activation without promoting CDK2 phosphorylation, by inducing a conformation change of the CDK2 T-loop that obstructs the substrate-binding cleft prior to kinase activation (PubMed:28666995). Mediates cell survival during the DNA damage process through activation of CDK2 (PubMed:12839962). [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RC210559L3