

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC210626L4V

HSPC142 (BABAM1) (NM_001033549) Human Tagged ORF Clone Lentiviral Particle

Product data:

| Product Type: | Lentiviral Particles |
|------------------------------|---|
| Product Name: | HSPC142 (BABAM1) (NM_001033549) Human Tagged ORF Clone Lentiviral Particle |
| Symbol: | HSPC142 |
| Synonyms: | C19orf62; HSPC142; MERIT40; NBA1 |
| Mammalian Cell Selection: | Puromycin |
| Vector: | pLenti-C-mGFP-P2A-Puro (PS100093) |
| Tag: | mGFP |
| ACCN: | NM_001033549 |
| ORF Size: | 987 bp |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(RC210626). |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u> |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| RefSeq: | <u>NM 001033549.1</u> |
| RefSeq Size: | 1505 bp |
| RefSeq ORF: | 990 bp |
| Locus ID: | 29086 |
| UniProt ID: | <u>Q9NWV8</u> |
| Cytogenetics: | 19p13.11 |
| MW: | 36.6 kDa |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

SORIGENE HSPC142 (BABA

HSPC142 (BABAM1) (NM_001033549) Human Tagged ORF Clone Lentiviral Particle – RC210626L4V

Gene Summary: Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX. In the BRCA1-A complex, it is required for the complex integrity and its localization at DSBs. Component of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin in various substrates (PubMed:24075985, PubMed:26195665). In these 2 complexes, it is probably required to maintain the stability of BABAM2 and help the 'Lys-63'-linked deubiguitinase activity mediated by BRCC3/BRCC36 component. The BRISC complex is required for normal mitotic spindle assembly and microtubule attachment to kinetochores via its role in deubiquitinating NUMA1 (PubMed:26195665). Plays a role in interferon signaling via its role in the deubiguitination of the interferon receptor IFNAR1; deubiguitination increases IFNAR1 activity by enhancing its stability and cell surface expression (PubMed:24075985). Downregulates the response to bacterial lipopolysaccharide (LPS) via its role in IFNAR1 deubiguitination (PubMed:24075985).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US