

Product datasheet for RC210714L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CRSP7 (MED26) (NM 004831) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CRSP7 (MED26) (NM_004831) Human Tagged ORF Clone Lentiviral Particle

Symbol:

CRSP7; CRSP70 Synonyms:

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag: NM 004831 ACCN: **ORF Size:** 1800 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC210714).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 004831.3

RefSeq Size: 3184 bp RefSeq ORF: 1803 bp Locus ID: 9441 **UniProt ID:** 095402 Cytogenetics: 19p13.11

Protein Families: Druggable Genome

MW: 65.4 kDa







Gene Summary:

The activation of gene transcription is a multistep process that is triggered by factors that recognize transcriptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e.g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors. [provided by RefSeq, Jul 2008]