

Product datasheet for RC210781L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Proteinase Activated Receptor 3 (F2RL2) (NM_004101) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Proteinase Activated Receptor 3 (F2RL2) (NM_004101) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Proteinase Activated Receptor 3

Synonyms: PAR-3; PAR3

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM_004101

ORF Size: 1122 bp

ORF Nucleotide

Sequence:

Domains:

The ORF insert of this clone is exactly the same as(RC210781).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 004101.2</u>

 RefSeq Size:
 3446 bp

 RefSeq ORF:
 1125 bp

 Locus ID:
 2151

 UniProt ID:
 000254

 Cytogenetics:
 5q13.3

7tm 1





Proteinase Activated Receptor 3 (F2RL2) (NM_004101) Human Tagged ORF Clone Lentiviral Particle – RC210781L1V

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Neuroactive ligand-receptor interaction

MW: 42.5 kDa

Gene Summary: This gene encodes a member of the protease-activated receptor (PAR) family which is a

subfamily of the seven transmembrane G protein-coupled cell surface receptor family. The encoded protein acts as a cofactor in the thrombin-mediated cleavage and activation of the protease-activated receptor family member PAR4. The encoded protein plays an essential role in hemostasis and thrombosis. Alternate splicing results in multiple transcript variants

that encode different isoforms. [provided by RefSeq, Feb 2012]