

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC210829L1V

## MSL2L1 (MSL2) (NM\_018133) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	MSL2L1 (MSL2) (NM_018133) Human Tagged ORF Clone Lentiviral Particle
Symbol:	MSL2L1
Synonyms:	MSL-2; MSL2L1; RNF184
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_018133
ORF Size:	1731 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC210829).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 018133.2</u>
RefSeq Size:	4696 bp
RefSeq ORF:	1734 bp
Locus ID:	55167
UniProt ID:	<u>Q9HCI7</u>
Cytogenetics:	3q22.3
Protein Families:	Druggable Genome
MW:	62.5 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Component of histone acetyltransferase complex responsible for the majority of histone H4<br/>acetylation at lysine 16 which is implicated in the formation of higher-order chromatin<br/>structure. Acts as an E3 ubiquitin ligase that promotes monoubiquitination of histone H2B at<br/>'Lys-35' (H2BK34Ub), but not that of H2A. This activity is greatly enhanced by<br/>heterodimerization with MSL1. H2B ubiquitination in turn stimulates histine H3 methylation<br/>at 'Lys-4' (H3K4me) and 'Lys-79' (H3K79me) and leads to gene activation, including that of<br/>HOXA9 and MEIS1.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US