

## Product datasheet for **RC210875L3V**

### LONP2 (NM\_031490) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Lentiviral Particles   |
| Product Name:             | LONP2 (NM_031490) Human Tagged ORF Clone Lentiviral Particle   |
| Symbol:                   | LONP2  |
| Synonyms:                 | LONP; LONPL; PLON; PSLON   |
| Mammalian Cell Selection: | Puromycin  |
| Vector:                   | pLenti-C-Myc-DDK-P2A-Puro (PS100092)   |
| Tag:                      | Myc-DDK  |
| ACCN:                     | NM_031490  |
| ORF Size:                 | 2556 bp  |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(RC210875).   |
| OTI Disclaimer:           | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a> |
| OTI Annotation:           | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| RefSeq:                   | <a href="#">NM_031490.2</a>  |
| RefSeq Size:              | 4342 bp  |
| RefSeq ORF:               | 2559 bp  |
| Locus ID:                 | 83752  |
| UniProt ID:               | <a href="#">Q86WA8</a>   |
| Cytogenetics:             | 16q12.1  |
| Domains:                  | Lon_C, AAA, AAA  |
| Protein Families:         | Druggable Genome, Protease   |



[View online »](#)

**MW:** 94.4 kDa

**Gene Summary:** In human, peroxisomes function primarily to catalyze fatty acid beta-oxidation and, as a by-product, produce hydrogen peroxide and superoxide. The protein encoded by this gene is an ATP-dependent protease that likely plays a role in maintaining overall peroxisome homeostasis as well as proteolytically degrading peroxisomal proteins damaged by oxidation. The protein has an N-terminal Lon N substrate recognition domain, an ATPase domain, a proteolytic domain, and, in some isoforms, a C-terminal peroxisome targeting sequence. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Jan 2017]