

#### OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for RC210884L4

#### NARG1 (NAA15) (NM\_057175) Human Tagged Lenti ORF Clone

### **Product data:**

Product Type:	Expression Plasmids
Product Name:	NARG1 (NAA15) (NM_057175) Human Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	NARG1
Synonyms:	Ga19; MRD50; NARG1; NAT1P; NATH; TBDN; TBDN100
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC210884).
<b>Restriction Sites:</b>	Sgfl-Mlul
Cloning Scheme:	
	Cloning sites used for ORF Shuttling:
	Sgf1         ORF         Mlu I            GCG ATC GCIC         ATG // NNÑ         ACG CGT



ACCN: ORF Size: NM\_057175 2598 bp




View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Section 2012 CRIGENE NARG1 (NAA15) (NM_057175) Human Tagged Lenti ORF Clone – RC210884L4	
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
RefSeq:	<u>NM 057175.3</u>
RefSeq Size:	5554 bp
RefSeq ORF:	2601 bp
Locus ID:	80155
UniProt ID:	<u>Q9BXJ9</u>
Cytogenetics:	4q31.1
Domains:	TPR
Protein Families:	Druggable Genome, Transcription Factors
MW:	101.1 kDa
Gene Summary:	N-alpha-acetylation is among the most common post-translational protein modifications in eukaryotic cells. This process involves the transfer of an acetyl group from acetyl-coenzyme A to the alpha-amino group on a nascent polypeptide and is essential for normal cell function. This gene encodes the auxillary subunit of the N-terminal acetyltransferase A (NatA) complex. [provided by RefSeq, Jan 2017]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

## **Product images:**



Circular map for RC210884L4

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US