

Product datasheet for RC210898L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SFRS9 (SRSF9) (NM_003769) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: SFRS9 (SRSF9) (NM_003769) Human Tagged ORF Clone Lentiviral Particle

Symbol: SFRS9

Synonyms: SFRS9; SRp30c

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_003769

ORF Size: 663 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC210898).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 003769.2

 RefSeq Size:
 1204 bp

 RefSeq ORF:
 666 bp

 Locus ID:
 8683

 UniProt ID:
 Q13242

 Cytogenetics:
 12q24.31

Domains: RRM

Protein Families: Druggable Genome







Protein Pathways: Spliceosome

MW: 25.5 kDa

Gene Summary: The protein encoded by this gene is a member of the serine/arginine (SR)-rich family of pre-

mRNA splicing factors, which constitute part of the spliceosome. Each of these factors contains an RNA recognition motif (RRM) for binding RNA and an RS domain for binding other proteins. The RS domain is rich in serine and arginine residues and facilitates interaction between different SR splicing factors. In addition to being critical for mRNA splicing, the SR

proteins have also been shown to be involved in mRNA export from the nucleus and in translation. Two pseudogenes, one on chromosome 15 and the other on chromosome 21,

have been found for this gene. [provided by RefSeq, Sep 2010]