

## Product datasheet for **RC210914L2V**

### AVPR V2 (AVPR2) (NM\_000054) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Lentiviral Particles   |
| Product Name:             | AVPR V2 (AVPR2) (NM_000054) Human Tagged ORF Clone Lentiviral Particle   |
| Symbol:                   | AVPR V2  |
| Synonyms:                 | ADHR; DI1; DIR; DIR3; NDI; NDI1; V2R   |
| Mammalian Cell Selection: | None   |
| Vector:                   | pLenti-C-mGFP (PS100071)   |
| Tag:                      | mGFP   |
| ACCN:                     | NM_000054  |
| ORF Size:                 | 1113 bp  |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(RC210914).   |
| OTI Disclaimer:           | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a> |
| OTI Annotation:           | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| RefSeq:                   | <a href="#">NM_000054.2</a>  |
| RefSeq Size:              | 1682 bp  |
| RefSeq ORF:               | 1116 bp  |
| Locus ID:                 | 554  |
| UniProt ID:               | <a href="#">P30518</a>   |
| Cytogenetics:             | Xq28   |
| Protein Families:         | Druggable Genome, GPCR, Transmembrane  |
| Protein Pathways:         | Neuroactive ligand-receptor interaction  |



[View online »](#)

MW: 40.3 kDa

**Gene Summary:** This gene encodes the vasopressin receptor, type 2, also known as the V2 receptor, which belongs to the seven-transmembrane-domain G protein-coupled receptor (GPCR) superfamily, and couples to Gs thus stimulating adenylate cyclase. The subfamily that includes the V2 receptor, the V1a and V1b vasopressin receptors, the oxytocin receptor, and isotocin and mesotocin receptors in non-mammals, is well conserved, though several members signal via other G proteins. All bind similar cyclic nonapeptide hormones. The V2 receptor is expressed in the kidney tubule, predominantly in the distal convoluted tubule and collecting ducts, where its primary property is to respond to the pituitary hormone arginine vasopressin (AVP) by stimulating mechanisms that concentrate the urine and maintain water homeostasis in the organism. When the function of this gene is lost, the disease Nephrogenic Diabetes Insipidus (NDI) results. The V2 receptor is also expressed outside the kidney although its tissue localization is uncertain. When these 'extrarenal receptors' are stimulated by infusion of a V2 selective agonist (dDAVP), a variety of clotting factors are released into the bloodstream. The physiologic importance of this property is not known - its absence does not appear to be detrimental in NDI patients. The gene expression has also been described in fetal lung tissue and lung cancer associated with alternative splicing. [provided by RefSeq, Jul 2008]