

## Product datasheet for RC211107L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## C1orf77 (CHTOP) (NM 015607) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** C1orf77 (CHTOP) (NM\_015607) Human Tagged ORF Clone Lentiviral Particle

Symbol: C1orf77

Synonyms: C1orf77; C10orf77; FL-SRAG; FOP; pp7704; SRAG; SRAG-3; SRAG-5

**Mammalian Cell** 

Selection:

ACCN:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 015607

Tag: Myc-DDK

ORF Size: 747 bp

**ORF Nucleotide** 

Tho C

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC211107).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 015607.2</u>

 RefSeq Size:
 2172 bp

 RefSeq ORF:
 747 bp

 Locus ID:
 26097

 UniProt ID:
 Q9Y3Y2

 Cytogenetics:
 1q21.3

 MW:
 26.5 kDa







## **Gene Summary:**

This gene encodes a small nuclear protein that is characterized by an arginine and glycine rich region. This protein may have an important role in the regulation of fetal globin gene expression and in the activation of estrogen-responsive genes. A recent study reported that this protein binds 5-hydroxymethylcytosine (5hmC) and associates with an arginine methyltransferase complex (methylosome), which promotes methylation of arginine 3 of histone H4 (H4R3) and activation of genes involved in glioblastomagenesis. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Nov 2015]