

Product datasheet for RC211151L1

HYAL4 (NM_012269) Human Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	HYAL4 (NM_012269) Human Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	HYAL4
Synonyms:	CSHY; HYAL-4
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC211151).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_012269
ORF Size:	1443 bp




[View online >](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_012269.1
RefSeq Size:	2411 bp
RefSeq ORF:	1446 bp
Locus ID:	23553
UniProt ID:	Q2M3T9
Cytogenetics:	7q31.32
Domains:	Glyco_hydro_56
Protein Pathways:	Glycosaminoglycan degradation, Metabolic pathways
MW:	54.3 kDa
Gene Summary:	This gene encodes a protein which is similar in structure to hyaluronidases but lacks hyaluronidase activity. The encoded protein acts as a chondroitin-sulfate-specific endo-beta-N-acetylgalactosaminidase; that is, it exhibits hydrolytic activity toward chondroitin sulfate chains and degrades them into oligosaccharides. Proteoglycans are formed by the covalent linkage of chondroitin sulfate chains to protein. Proteoglycans are ubiquitous components of the extracellular matrix of connective tissues and are also found at the surface of many cell types where they participate in a variety of cellular processes such as cell proliferation, differentiation, migration, cell-cell recognition, extracellular matrix deposition, and tissue morphogenesis. The expression of this gene is highest in testes and placenta. [provided by RefSeq, Apr 2019]

Product images:



Circular map for RC211151L1



Double digestion of RC211151L1 using SgfI and MluI