

## Product datasheet for RC211228L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## LILRB3 (NM 006864) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** LILRB3 (NM 006864) Human Tagged ORF Clone Lentiviral Particle

Symbol:

CD85A; HL9; ILT-5; ILT5; LILRA6; LIR-3; LIR3; PIR-B; PIRB Synonyms:

**Mammalian Cell** 

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag: NM 006864 ACCN:

**ORF Size:** 1893 bp

**ORF Nucleotide** 

Sequence: OTI Disclaimer: The ORF insert of this clone is exactly the same as(RC211228).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 006864.2, NP 006855.1

RefSeq Size: 2840 bp RefSeq ORF: 1896 bp Locus ID: 11025 **Cytogenetics:** 19q13.42 Domains: ig, IGc2, IG

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** B cell receptor signaling pathway





ORIGENE

**MW:** 69.4 kDa

**Gene Summary:** 

This gene is a member of the leukocyte immunoglobulin-like receptor (LIR) family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]