

## Product datasheet for RC211253L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TNFRSF4 (NM 003327) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** TNFRSF4 (NM\_003327) Human Tagged ORF Clone Lentiviral Particle

Symbol:

ACT35; CD134; IMD16; OX40; TXGP1L Synonyms:

**Mammalian Cell** 

Selection:

ACCN:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Myc-DDK Tag: NM 003327

**ORF Size:** 831 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC211253).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 003327.2

RefSeq Size: 1120 bp RefSeq ORF: 834 bp Locus ID: 7293 **UniProt ID:** P43489 Cytogenetics: 1p36.33 **Domains: TNFR** 

**Protein Families:** Transmembrane





## TNFRSF4 (NM\_003327) Human Tagged ORF Clone Lentiviral Particle - RC211253L3V

**Protein Pathways:** Cytokine-cytokine receptor interaction

MW: 29.3 kDa

**Gene Summary:** The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor

has been shown to activate NF-kappaB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that this receptor promotes the expression of

apoptosis inhibitors BCL2 and BCL2IL1/BCL2-XL, and thus suppresses apoptosis. The

knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul 2008]