

Product datasheet for RC211260L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cyclin B2 (CCNB2) (NM_004701) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Cyclin B2 (CCNB2) (NM_004701) Human Tagged ORF Clone Lentiviral Particle

Symbol: Cyclin B2
Synonyms: HsT17299
Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 004701

ORF Size: 1194 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC211260).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 004701.2

 RefSeq Size:
 1566 bp

 RefSeq ORF:
 1197 bp

 Locus ID:
 9133

 UniProt ID:
 095067

 Cytogenetics:
 15q22.2

Domains: cyclin_C, CYCLIN, cyclin

Protein Families: Druggable Genome, Stem cell - Pluripotency





Cyclin B2 (CCNB2) (NM_004701) Human Tagged ORF Clone Lentiviral Particle - RC211260L3V

Protein Pathways: Cell cycle, Oocyte meiosis, p53 signaling pathway, Progesterone-mediated oocyte maturation

MW: 45.3 kDa

Gene Summary: Cyclin B2 is a member of the cyclin family, specifically the B-type cyclins. The B-type cyclins, B1

and B2, associate with p34cdc2 and are essential components of the cell cycle regulatory machinery. B1 and B2 differ in their subcellular localization. Cyclin B1 co-localizes with microtubules, whereas cyclin B2 is primarily associated with the Golgi region. Cyclin B2 also binds to transforming growth factor beta RII and thus cyclin B2/cdc2 may play a key role in transforming growth factor beta-mediated cell cycle control. [provided by RefSeq, Jul 2008]