

Product datasheet for **RC211306**

NKG2C (KLRC2) (NM_002260) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	NKG2C (KLRC2) (NM_002260) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	NKG2C
Synonyms:	CD159c; NKG2-C; NKG2C
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC211306 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGAGTAAACAAAGAGGAACCTTCTCAGAAGTGAGTCTGGCCCAGGACCCAAAGCGGCAGCAAAGGAAAC
CTAAAGGCAATAAAAGCTCCATTCAGGAACCGAACAGGAAATATTCCAAGTAGAATTAATCTTCAAAA
TCCTTCCCTGAATCATCAAGGGATTGATAAAATATATGACTGCCAAGTTTACTGCCACCTCCAGAGAAG
CTCACTGCCGAGGTCCTAGGAATCATTTCATTGCTCTGATGGCCACTGTGTTAAAAACAATAGTTCCTTA
TTCCTTTCCTGGAGCAGAACAAATCTTCCCGAATACAAGAACGCAGAAAGCACGTCATTGTGGCCATTG
TCCTGAGGAGTGGATTACATATTCCAACAGTTGTTATTACATTGGTAAGGAAAGAAGAAGAACTGGGAAGAG
AGTTTGCTGGCCCTGACTTCGAAGAACTCCAGTCTGCTTTCTATAGATAATGAAGAAGAAATGAAATTTT
TGGCCAGCATTTTACCTTCTCATGGATTGGTGTGTTTCGTAACAGCAGTCATCATCCATGGGTGACAAT
AAATGGTTTGGCTTTCAAACATAAGATAAAAGACTCAGATAATGCTGAAGTAACTGTGCAGTGCTACAA
GTAAATCGACTTAAATCAGCCCAGTGTGGATCTCAATGATATATCATTGTAAGCATAAGCTT

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC211306 protein sequence
Red=Cloning site Green=Tags(s)

MSKQRGTFSEVSLAQDPKRQQRKPKGNKSSISGTEQEIFQVELNLQNPSLNHQGIDKIYDCQGLLPPEK
 LTAEVLGIICIVLMATVLKTIIVLPFLEQNSSPNTRTQKARHCGHCPEEWITYSNSCYIIGKERRTWE
 SLLACTSKNSSLLSIDNEEEMKFLASILPSSWIGVFRNSSHPWVTINGLAFKHKIKDSDNAELNCAVLQ
 VNRLKSAQCGSSMIYHCKHKL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6369_f07.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_002260

ORF Size: 693 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.


Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_002260.3 , NP_002251.2
RefSeq Size:	1223 bp
RefSeq ORF:	696 bp
Locus ID:	3822
UniProt ID:	P26717
Cytogenetics:	12p13.2
Protein Families:	Transmembrane
Protein Pathways:	Antigen processing and presentation, Natural killer cell mediated cytotoxicity
MW:	26.1 kDa
Gene Summary:	<p>Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. The group, designated KLRC (NKG2) are expressed primarily in natural killer (NK) cells and encodes a family of transmembrane proteins characterized by a type II membrane orientation (extracellular C terminus) and the presence of a C-type lectin domain. The KLRC (NKG2) gene family is located within the NK complex, a region that contains several C-type lectin genes preferentially expressed on NK cells. KLRC2 alternative splice variants have been described but their full-length nature has not been determined. [provided by RefSeq, Jul 2008]</p>

Product images:



Circular map for RC211306



Western blot validation of overexpression lysate (Cat# [LY419440]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC211306 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).