

Product datasheet for RC211459L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

FLT3 (NM 004119) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: FLT3 (NM_004119) Human Tagged ORF Clone Lentiviral Particle

Symbol:

CD135; FLK-2; FLK2; STK1 Synonyms:

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK NM 004119 ACCN: **ORF Size:** 2979 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC211459).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 004119.1

RefSeq Size: 3475 bp RefSeq ORF: 2982 bp Locus ID: 2322 **UniProt ID:** P36888

Cytogenetics: 13q12.2

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Protein Kinase, Transmembrane



FLT3 (NM_004119) Human Tagged ORF Clone Lentiviral Particle - RC211459L3V

Protein Pathways: Acute myeloid leukemia, Cytokine-cytokine receptor interaction, Hematopoietic cell lineage,

Pathways in cancer

MW: 112.7 kDa

Gene Summary: This gene encodes a class III receptor tyrosine kinase that regulates hematopoiesis. This

receptor is activated by binding of the fms-related tyrosine kinase 3 ligand to the extracellular

domain, which induces homodimer formation in the plasma membrane leading to autophosphorylation of the receptor. The activated receptor kinase subsequently

phosphorylates and activates multiple cytoplasmic effector molecules in pathways involved in

apoptosis, proliferation, and differentiation of hematopoietic cells in bone marrow.

Mutations that result in the constitutive activation of this receptor result in acute myeloid leukemia and acute lymphoblastic leukemia. [provided by RefSeq, Jan 2015]