

## Product datasheet for RC211487L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MRGX1 (MRGPRX1) (NM 147199) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: MRGX1 (MRGPRX1) (NM\_147199) Human Tagged ORF Clone Lentiviral Particle

Symbol: MRGX1

Synonyms: GPCR; MGRG2; MRGX1; SNSR4

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 147199

ORF Size: 966 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC211487).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 147199.3, NP 671732.3

RefSeq Size: 1190 bp
RefSeq ORF: 969 bp
Locus ID: 259249
UniProt ID: Q96LB2

Cytogenetics: 11

**Protein Families:** Druggable Genome, GPCR, Transmembrane

MW: 36.3 kDa





## **Gene Summary:**

Orphan receptor. Probably involved in the function of nociceptive neurons. May regulate nociceptor function and/or development, including the sensation or modulation of pain. Potently activated by enkephalins including BAM22 (bovine adrenal medulla peptide 22) and BAM (8-22)(PubMed:26582731). BAM22 is the most potent compound and evoked a large and dose-dependent release of intracellular calcium in stably transfected cells. G(alpha)q proteins are involved in the calcium-signaling pathway. Activated by the antimalarial drug, chloroquine. May mediate chloroquine-induced itch, in a histamine-independent manner. [UniProtKB/Swiss-Prot Function]