

## Product datasheet for RC211582L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## BTK (NM\_000061) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** BTK (NM\_000061) Human Tagged ORF Clone Lentiviral Particle

Symbol: BTK

Synonyms: AGMX1; AT; ATK; BPK; IGHD3; IMD1; PSCTK1; XLA

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM\_000061

 ORF Size:
 1977 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC211582).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 000061.1

RefSeq Size:2591 bpRefSeq ORF:1980 bp

Locus ID: 695

UniProt ID: Q06187

Cytogenetics: Xq22.1

**Domains:** pkinase, SH2, TyrKc, SH3, BTK, PH, S\_TKc

**Protein Families:** Druggable Genome, Protein Kinase



## BTK (NM\_000061) Human Tagged ORF Clone Lentiviral Particle - RC211582L1V

**Protein Pathways:** B cell receptor signaling pathway, Fc epsilon RI signaling pathway, Primary immunodeficiency

**MW:** 76.1 kDa

**Gene Summary:** The protein encoded by this gene plays a crucial role in B-cell development. Mutations in this

gene cause X-linked agammaglobulinemia type 1, which is an immunodeficiency

characterized by the failure to produce mature B lymphocytes, and associated with a failure of Ig heavy chain rearrangement. Alternative splicing results in multiple transcript variants

encoding different isoforms. [provided by RefSeq, Dec 2013]