

Product datasheet for RC211599L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

cIAP2 (BIRC3) (NM_182962) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: cIAP2 (BIRC3) (NM_182962) Human Tagged ORF Clone Lentiviral Particle

Symbol: cIAP2

Synonyms: AIP1; API2; c-IAP2; CIAP2; HAIP1; HIAP1; IAP-1; MALT2; MIHC; RNF49

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_182962 **ORF Size:** 1812 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC211599).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 182962.1

 RefSeq Size:
 4372 bp

 RefSeq ORF:
 1815 bp

 Locus ID:
 330

 UniProt ID:
 Q13489

Cytogenetics: 11q22.2

Protein Families: Druggable Genome





cIAP2 (BIRC3) (NM_182962) Human Tagged ORF Clone Lentiviral Particle - RC211599L4V

Protein Pathways: Apoptosis, Focal adhesion, NOD-like receptor signaling pathway, Pathways in cancer, Small

cell lung cancer, Ubiquitin mediated proteolysis

MW: 68.4 kDa

Gene Summary: This gene encodes a member of the IAP family of proteins that inhibit apoptosis by binding to

tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. The encoded protein inhibits apoptosis induced by serum deprivation but does not affect apoptosis resulting from exposure to menadione, a potent inducer of free radicals. It contains 3 baculovirus IAP repeats and a ring finger domain. Transcript variants encoding the same isoform have been identified. [provided by RefSeq,

Aug 2011]