

## Product datasheet for RC211635L2

### BCR (NM\_004327) Human Tagged Lenti ORF Clone

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Expression Plasmids  |
| Product Name:             | BCR (NM_004327) Human Tagged Lenti ORF Clone                   |
| Tag:                      | mGFP   |
| Symbol:                   | BCR  |
| Synonyms:                 | ALL; BCR1; CML; D22S11; D22S662; PHL                           |
| Mammalian Cell Selection: | None   |
| Vector:                   | pLenti-C-mGFP (PS100071)                                       |
| E. coli Selection:        | Chloramphenicol (34 ug/mL)                                     |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(RC211635). |
| Restriction Sites:        | SgfI-MluI  |
| Cloning Scheme:           |  |


Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.



[View online »](#)

**Plasmid Map:**


**ACCN:** NM\_004327

**ORF Size:** 3813 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_004327.3](#)

**RefSeq Size:** 4739 bp

**RefSeq ORF:** 3816 bp

|                          |  |
|--------------------------|--|
| <b>Locus ID:</b>         | 613  |
| <b>UniProt ID:</b>       | <a href="#">P11274</a>                       |
| <b>Cytogenetics:</b>     | 22q11.23                                     |
| <b>Domains:</b>          | C2, RhoGAP, RhoGEF, PH                       |
| <b>Protein Families:</b> | Druggable Genome, Protein Kinase             |
| <b>Protein Pathways:</b> | Chronic myeloid leukemia, Pathways in cancer |
| <b>MW:</b>               | 142.6 kDa                                    |

**Gene Summary:** A reciprocal translocation between chromosomes 22 and 9 produces the Philadelphia chromosome, which is often found in patients with chronic myelogenous leukemia. The chromosome 22 breakpoint for this translocation is located within the BCR gene. The translocation produces a fusion protein which is encoded by sequence from both BCR and ABL, the gene at the chromosome 9 breakpoint. Although the BCR-ABL fusion protein has been extensively studied, the function of the normal BCR gene product is not clear. The unregulated tyrosine kinase activity of BCR-ABL1 contributes to the immortality of leukaemic cells. The BCR protein has serine/threonine kinase activity and is a GTPase-activating protein for p21rac and other kinases. Two transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jan 2020]