

## Product datasheet for RC211673L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## DEP1 (PTPRJ) (NM 001098503) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: DEP1 (PTPR)) (NM 001098503) Human Tagged ORF Clone Lentiviral Particle

Symbol: DEP1

Synonyms: CD148; DEP1; HPTPeta; R-PTP-ETA; SCC1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

**ACCN:** NM\_001098503

ORF Size: 1617 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC211673).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001098503.1</u>

 RefSeq Size:
 3193 bp

 RefSeq ORF:
 1620 bp

 Locus ID:
 5795

 UniProt ID:
 Q12913

Cytogenetics: 11p11.2

**Protein Families:** Druggable Genome, Phosphatase, Transmembrane

**Protein Pathways:** Adherens junction





**MW:** 57.19 kDa

**Gene Summary:** 

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes, including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP possesses an extracellular region containing five fibronectin type III repeats, a single transmembrane region, and a single intracytoplasmic catalytic domain, and thus represents a receptor-type PTP. This protein is present in all hematopoietic lineages, and was shown to negatively regulate T cell receptor signaling possibly through interfering with the phosphorylation of Phospholipase C Gamma 1 and Linker for Activation of T Cells. This protein can also dephosphorylate the PDGF beta receptor, and may be involved in UV-induced signal transduction. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]