

## Product datasheet for RC211740L3

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### PDGF Receptor alpha (PDGFRA) (NM\_006206) Human Tagged Lenti ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

Product Name: PDGF Receptor alpha (PDGFRA) (NM\_006206) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: PDGF Receptor alpha

Synonyms: CD140A; PDGFR-2; PDGFR2

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF


Sequence:

The ORF insert of this clone is exactly the same as(RC211740).

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF.

**ACCN:** NM\_006206

ORF Size: 3267 bp



#### PDGF Receptor alpha (PDGFRA) (NM\_006206) Human Tagged Lenti ORF Clone - RC211740L3

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 006206.3</u>, <u>NP 006197.1</u>

4q12

 RefSeq Size:
 6405 bp

 RefSeq ORF:
 3270 bp

 Locus ID:
 5156

 UniProt ID:
 P16234

Cytogenetics:

**Domains:** pkinase, TyrKc, S TKc, ig, IGc2, IG

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Protein Kinase, Transmembrane

**Protein Pathways:** Calcium signaling pathway, Colorectal cancer, Cytokine-cytokine receptor interaction,

Endocytosis, Focal adhesion, Gap junction, Glioma, MAPK signaling pathway, Melanoma,

Pathways in cancer, Prostate cancer, Regulation of actin cytoskeleton

MW: 122.67 kDa


**Gene Summary:** This gene encodes a cell surface tyrosine kinase receptor for members of the platelet-derived

growth factor family. These growth factors are mitogens for cells of mesenchymal origin. The identity of the growth factor bound to a receptor monomer determines whether the functional receptor is a homodimer or a heterodimer, composed of both platelet-derived growth factor receptor alpha and beta polypeptides. Studies suggest that this gene plays a role in organ development, wound healing, and tumor progression. Mutations in this gene have been associated with idiopathic hypereosinophilic syndrome, somatic and familial gastrointestinal stromal tumors, and a variety of other cancers. [provided by RefSeq, Mar

2012]



# **Product images:**



Circular map for RC211740L3