

## Product datasheet for RC211741L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IL5RA (NM\_000564) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** IL5RA (NM\_000564) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL5RA

Synonyms: CD125; CDw125; HSIL5R3; IL5R

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 000564

ORF Size: 1260 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC211741).

Sequence:

Cytogenetics:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 000564.2

 RefSeq Size:
 2367 bp

 RefSeq ORF:
 1263 bp

 Locus ID:
 3568

 UniProt ID:
 Q01344

**Protein Families:** Druggable Genome, Transmembrane

3p26.2





## IL5RA (NM\_000564) Human Tagged ORF Clone Lentiviral Particle - RC211741L3V

**Protein Pathways:** Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, Jak-STAT signaling

pathway

MW: 47.5 kDa

**Gene Summary:** The protein encoded by this gene is an interleukin 5 specific subunit of a heterodimeric

cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3), colony stimulating factor 2 (CSF2/GM-CSF), and interleukin 5 (IL5). The binding of this protein to IL5 depends on the beta subunit. The beta subunit is activated by the ligand binding, and is required for the biological activities of IL5. This protein has been found to interact with syndecan binding protein (syntenin), which is required for IL5 mediated activation of the transcription factor SOX4. Several alternatively spliced transcript variants encoding four distinct isoforms have

been reported. [provided by RefSeq, Jul 2011]