

Product datasheet for RC211766L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Collagen III (COL3A1) (NM 000090) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Collagen III (COL3A1) (NM_000090) Human Tagged ORF Clone Lentiviral Particle

Symbol: Collagen III

Synonyms: EDS4A; EDSVASC; PMGEDSV

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK ACCN: NM_000090

ORF Size: 4398 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC211766).

Sequence:

Cytogenetics:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000090.2

 RefSeq Size:
 5489 bp

 RefSeq ORF:
 4401 bp

 Locus ID:
 1281

 UniProt ID:
 P02461

Domains: COLFI, VWC, Collagen

2q32.2

Protein Families: Druggable Genome





Collagen III (COL3A1) (NM_000090) Human Tagged ORF Clone Lentiviral Particle - RC211766L3V

Protein Pathways: ECM-receptor interaction, Focal adhesion

MW: 138.6 kDa

Gene Summary: This gene encodes the pro-alpha1 chains of type III collagen, a fibrillar collagen that is found

in extensible connective tissues such as skin, lung, uterus, intestine and the vascular system, frequently in association with type I collagen. Mutations in this gene are associated with Ehlers-Danlos syndrome types IV, and with aortic and arterial aneurysms. Two transcripts, resulting from the use of alternate polyadenylation signals, have been identified for this gene.

[provided by R. Dalgleish, Feb 2008]