

Product datasheet for RC211784

p16INK4A (CDKN2A) (NM_058195) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Tag:	Myc-DDK
Symbol:	p16INK4A
Synonyms:	ARF; CDK4I; CDKN2; CMM2; INK4; INK4A; MLM; MTS-1; MTS1; P14; P14ARF; P16; P16-INK4A; P16INK4; P16INK4A; P19; P19ARF; TP16
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC211784 representing NM_058195 Red=Cloning site Blue=ORF Green=Tags(s)

TTTGTGAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCGCGATCGCC

ATGGGCAGGGGGCGGTGCGTGGGTCCCAGTCTGCAGTTAAGGGGGCAGGAGTGGCGCTGCTCACCTCTGG
 TGCCAAAGGGCGGCGCAGCGGCTGCCGAGCTCGGCCCTGGAGGCGGCGAGAACATGGTGCGCAGGTTCTT
 GGTGACCTCCGATTCCGCGCGCGTGC GGCCCGCGAGTGAGGGTTTCGTGGTTCACATCCCGCGG
 CTCACGGGGAGTGGGCAGCGCCAGGGGCGCCGCGCTGTGGCCCTCGTGCTGATGCTACTGAGGAGCC
 AGCGTCTAGGGCAGCAGCCGCTTCTAGAAGACCAGGTCATGATGATGGGCAGCGCCGAGTGGCGGAGC
 TGCTGCTGCTCCACGGCGCGGAGCCCACTGCGCCGACCCGCCACTCTACCCGACCCGTGCACGACGC
 TGCCCGGGAGGGCTTCTGGACACGCTGGTGGTGTGCTGCACCGGGCGGGGCGCGCTGGACGTGC GCGAT
 GCCTGGGGCCGTCTGCCCGTGGACCTGGC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence:	>RC211784 representing NM_058195 Red=Cloning site Green=Tags(s)
-------------------	--

MGRGRCVGPSLQLRGQEWRCSP LVPKGGAAAAELGPGGGEMVRRFLVTLRIRACGP PPRVRFV VVHIPR
 LTGEWAAPGAPAAVALV LMLLRSQLGQQPLPRRPGHDDGQRP SGAAAAAPRRGAQLRRPRHSHPTRARR
 CPGGLPGHAGGAAPGRGAAGRARCLGPSARGPG


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms:	https://cdn.origene.com/chromatograms/mg3950_f09.zip
----------------	---


[View online »](#)

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_058195

ORF Size: 519 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.


RefSeq: [NM_058195.2](#), [NP_478102.1](#)

RefSeq Size: 1154 bp


RefSeq ORF: 399 bp

Locus ID:	1029
UniProt ID:	Q8N726
Cytogenetics:	9p21.3
Protein Families:	Druggable Genome
Protein Pathways:	Bladder cancer, Cell cycle, Chronic myeloid leukemia, Glioma, Melanoma, Non-small cell lung cancer, p53 signaling pathway, Pancreatic cancer, Pathways in cancer
MW:	17.8 kDa
Gene Summary:	<p>This gene generates several transcript variants which differ in their first exons. At least three alternatively spliced variants encoding distinct proteins have been reported, two of which encode structurally related isoforms known to function as inhibitors of CDK4 kinase. The remaining transcript includes an alternate first exon located 20 Kb upstream of the remainder of the gene; this transcript contains an alternate open reading frame (ARF) that specifies a protein which is structurally unrelated to the products of the other variants. This ARF product functions as a stabilizer of the tumor suppressor protein p53 as it can interact with, and sequester, the E3 ubiquitin-protein ligase MDM2, a protein responsible for the degradation of p53. In spite of the structural and functional differences, the CDK inhibitor isoforms and the ARF product encoded by this gene, through the regulatory roles of CDK4 and p53 in cell cycle G1 progression, share a common functionality in cell cycle G1 control. This gene is frequently mutated or deleted in a wide variety of tumors, and is known to be an important tumor suppressor gene. [provided by RefSeq, Sep 2012]</p>


Product images:



Circular map for RC211784



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY CDKN2A (Cat# RC211784, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CDKN2A (Cat# [TA802246]). Positive lysates [LY429923] (100ug) and [LC429923] (20ug) can be purchased separately from OriGene.



Western blot validation of overexpression lysate (Cat# [LY409229]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC211784 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).