

## Product datasheet for RC211795L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TAK1 (MAP3K7) (NM\_145331) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: TAK1 (MAP3K7) (NM 145331) Human Tagged ORF Clone Lentiviral Particle

Symbol: TAK1

**Synonyms:** CSCF; FMD2; MEKK7; TAK1; TGF1a

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM\_145331

 ORF Size:
 1818 bp

**ORF Nucleotide** 

- - -

Sequence:

**UniProt ID:** 

The ORF insert of this clone is exactly the same as(RC211795).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 145331.1</u>

 RefSeq Size:
 2850 bp

 RefSeq ORF:
 1821 bp

 Locus ID:
 6885

**Cytogenetics:** 6q15

**Protein Families:** Druggable Genome, Protein Kinase

O43318





## TAK1 (MAP3K7) (NM\_145331) Human Tagged ORF Clone Lentiviral Particle - RC211795L1V

Protein Pathways: Adherens junction, MAPK signaling pathway, NOD-like receptor signaling pathway, RIG-I-like

receptor signaling pathway, T cell receptor signaling pathway, Toll-like receptor signaling

pathway, Wnt signaling pathway

**MW:** 67 kDa

**Gene Summary:** The protein encoded by this gene is a member of the serine/threonine protein kinase family.

This kinase mediates the signaling transduction induced by TGF beta and morphogenetic protein (BMP), and controls a variety of cell functions including transcription regulation and

apoptosis. In response to IL-1, this protein forms a kinase complex including TRAF6,

MAP3K7P1/TAB1 and MAP3K7P2/TAB2; this complex is required for the activation of nuclear factor kappa B. This kinase can also activate MAPK8/JNK, MAP2K4/MKK4, and thus plays a role in the cell response to environmental stresses. Four alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]