

## Product datasheet for RC211813L3V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# PRC1 (NM\_199413) Human Tagged ORF Clone Lentiviral Particle

#### **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** PRC1 (NM\_199413) Human Tagged ORF Clone Lentiviral Particle

Symbol: PRC1 Synonyms: ASE1

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_199413

 ORF Size:
 1818 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC211813).

•

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 199413.1

RefSeq Size: 3091 bp
RefSeq ORF: 1821 bp
Locus ID: 9055
Cytogenetics: 15q26.1

**MW:** 70.1 kDa







#### **Gene Summary:**

This gene encodes a protein that is involved in cytokinesis. The protein is present at high levels during the S and G2/M phases of mitosis but its levels drop dramatically when the cell exits mitosis and enters the G1 phase. It is located in the nucleus during interphase, becomes associated with mitotic spindles in a highly dynamic manner during mitosis, and localizes to the cell mid-body during cytokinesis. This protein has been shown to be a substrate of several cyclin-dependent kinases (CDKs). It is necessary for polarizing parallel microtubules and concentrating the factors responsible for contractile ring assembly. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2012]