

Product datasheet for RC211819L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

iNOS (NOS2) (NM_000625) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: iNOS (NOS2) (NM_000625) Human Tagged ORF Clone Lentiviral Particle

Symbol: iNOS

Synonyms: HEP-NOS; INOS; NOS; NOS2A

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_000625 **ORF Size:** 3459 bp

ORF Nucleotide

OTI Disclaimer:

3 133 SP

Sequence:

The ORF insert of this clone is exactly the same as(RC211819).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000625.3

 RefSeq Size:
 4221 bp

 RefSeq ORF:
 3462 bp

 Locus ID:
 4843

 UniProt ID:
 P35228

 Cytogenetics:
 17q11.2

Protein Families: Druggable Genome





iNOS (NOS2) (NM_000625) Human Tagged ORF Clone Lentiviral Particle - RC211819L4V

Protein Pathways: Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Arginine and proline metabolism,

Calcium signaling pathway, Long-term depression, Metabolic pathways, Pathways in cancer,

Small cell lung cancer

MW: 130.9 kDa

Gene Summary: Nitric oxide is a reactive free radical which acts as a biologic mediator in several processes,

including neurotransmission and antimicrobial and antitumoral activities. This gene encodes a nitric oxide synthase which is expressed in liver and is inducible by a combination of lipopolysaccharide and certain cytokines. Three related pseudogenes are located within the

Smith-Magenis syndrome region on chromosome 17. [provided by RefSeq, Jul 2008]