

## Product datasheet for RC211922L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Inhibin beta C chain (INHBC) (NM\_005538) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Inhibin beta C chain (INHBC) (NM\_005538) Human Tagged ORF Clone Lentiviral Particle

Symbol: Inhibin beta C chain

Synonyms: IHBC

Mammalian Cell None

Selection:

Vector:

pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

**ACCN:** NM\_005538

ORF Size: 1056 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC211922).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 005538.2

RefSeq Size: 1990 bp
RefSeq ORF: 1059 bp
Locus ID: 3626

UniProt ID: P55103

Cytogenetics: 12q13.3

**Domains:** TGF-beta

**Protein Families:** Druggable Genome, Secreted Protein





## Inhibin beta C chain (INHBC) (NM\_005538) Human Tagged ORF Clone Lentiviral Particle – RC211922L1V

**Protein Pathways:** Cytokine-cytokine receptor interaction, TGF-beta signaling pathway

MW: 38.21 kDa

**Gene Summary:** This gene encodes a member of the TGF-beta (transforming growth factor-beta) superfamily

of proteins. The encoded preproprotein is proteolytically processed to generate a subunit of homodimeric and heterodimeric activin complexes. The heterodimeric complex may function in the inhibition of activin A signaling. Transgenic mice overexpressing this gene exhibit

defects in testis, liver and prostate. [provided by RefSeq, Aug 2016]