

Product datasheet for RC211922L4

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Inhibin beta C chain (INHBC) (NM_005538) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Inhibin beta C chain (INHBC) (NM_005538) Human Tagged Lenti ORF Clone

Tag: mGFP

Symbol: Inhibin beta C chain

Synonyms: IHBC

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clo


Sequence:

The ORF insert of this clone is exactly the same as(RC211922).

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_005538

ORF Size: 1056 bp



Inhibin beta C chain (INHBC) (NM_005538) Human Tagged Lenti ORF Clone - RC211922L4

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 005538.2</u>

 RefSeq Size:
 1990 bp

 RefSeq ORF:
 1059 bp

 Locus ID:
 3626

 UniProt ID:
 P55103

Cytogenetics: 12q13.3

Domains: TGF-beta

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Cytokine-cytokine receptor interaction, TGF-beta signaling pathway

MW: 38.21 kDa


Gene Summary: This gene encodes a member of the TGF-beta (transforming growth factor-beta) superfamily

of proteins. The encoded preproprotein is proteolytically processed to generate a subunit of homodimeric and heterodimeric activin complexes. The heterodimeric complex may function in the inhibition of activin A signaling. Transgenic mice overexpressing this gene exhibit

defects in testis, liver and prostate. [provided by RefSeq, Aug 2016]



Product images:



Circular map for RC211922L4