

Product datasheet for RC211922L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Inhibin beta C chain (INHBC) (NM 005538) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Inhibin beta C chain (INHBC) (NM_005538) Human Tagged ORF Clone Lentiviral Particle

Symbol: Inhibin beta C chain

Synonyms: IHBC

Mammalian Cell

Puromycin

Selection:

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_005538 **ORF Size:** 1056 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC211922).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 005538.2

 RefSeq Size:
 1990 bp

 RefSeq ORF:
 1059 bp

 Locus ID:
 3626

 UniProt ID:
 P55103

 Cytogenetics:
 12q13.3

 Domains:
 TGF-beta

Protein Families: Druggable Genome, Secreted Protein





Inhibin beta C chain (INHBC) (NM_005538) Human Tagged ORF Clone Lentiviral Particle – RC211922L4V

Protein Pathways: Cytokine-cytokine receptor interaction, TGF-beta signaling pathway

MW: 38.21 kDa

Gene Summary: This gene encodes a member of the TGF-beta (transforming growth factor-beta) superfamily

of proteins. The encoded preproprotein is proteolytically processed to generate a subunit of homodimeric and heterodimeric activin complexes. The heterodimeric complex may function in the inhibition of activin A signaling. Transgenic mice overexpressing this gene exhibit

defects in testis, liver and prostate. [provided by RefSeq, Aug 2016]