

Product datasheet for RC212044L2

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Eotaxin 3 (CCL26) (NM_006072) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Eotaxin 3 (CCL26) (NM 006072) Human Tagged Lenti ORF Clone

Tag: mGFP

Symbol: Eotaxin 3

Synonyms: IMAC; MIP-4a; MIP-4alpha; SCYA26; TSC-1

Mammalian Cell None

Selection:

Vector:pLenti-C-mGFP (PS100071)E. coli Selection:Chloramphenicol (34 ug/mL)


ORF Nucleotide The ORF insert of this clone is exactly the same as(RC212044).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_006072

ORF Size: 282 bp



Eotaxin 3 (CCL26) (NM_006072) Human Tagged Lenti ORF Clone - RC212044L2

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 006072.4</u>, <u>NP 006063.1</u>

 RefSeq Size:
 562 bp

 RefSeq ORF:
 285 bp

 Locus ID:
 10344

 UniProt ID:
 Q9Y258

 Cytogenetics:
 7q11.23

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Chemokine signaling pathway, Cytokine-cytokine receptor interaction

MW: 10.65 kDa


Gene Summary: This gene is one of two Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 7.

Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for normal peripheral blood eosinophils and basophils. This protein also has antimicrobial activity, displaying an antibacterial effect on S. pneumoniae, S. aureus, Non-typeable H. influenzae, and P. aeruginosa. The product of this gene is one of three related chemokines that specifically activate chemokine receptor CCR3. This chemokine may contribute to the eosinophil

accumulation in atopic diseases. [provided by RefSeq, Jul 2020]



Product images:



Circular map for RC212044L2