

## Product datasheet for RC212055L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## BCL2 (NM\_000657) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** BCL2 (NM\_000657) Human Tagged ORF Clone Lentiviral Particle

Symbol: BCL2

Synonyms: Bcl-2; PPP1R50

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_000657

ORF Size: 615 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC212055).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 000657.2

 RefSeq Size:
 1207 bp

 RefSeq ORF:
 618 bp

 Locus ID:
 596

 UniProt ID:
 P10415

Cytogenetics: 18q21.33

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Stem cell - Pluripotency, Transmembrane





## BCL2 (NM\_000657) Human Tagged ORF Clone Lentiviral Particle - RC212055L2V

**Protein Pathways:** Amyotrophic lateral sclerosis (ALS), Apoptosis, Colorectal cancer, Focal adhesion,

Neurotrophin signaling pathway, Pathways in cancer, Prostate cancer, Small cell lung cancer

MW: 22.2 kDa

**Gene Summary:** This gene encodes an integral outer mitochondrial membrane protein that blocks the

apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Alternative splicing results in multiple transcript variants. [provided by

RefSeq, Feb 2016]