

Product datasheet for RC212084L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL9R (NM_002186) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: IL9R (NM_002186) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL9R

Synonyms: CD129; IL-9R

Mammalian Cell None

Selection:

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_002186 **ORF Size:** 1563 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC212084).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 002186.2</u>

RefSeq Size: 2171 bp
RefSeq ORF: 1566 bp
Locus ID: 3581
UniProt ID: Q01113

Cytogenetics: Xq28 and Yq12

Protein Families: Druggable Genome, Secreted Protein, Transmembrane





IL9R (NM_002186) Human Tagged ORF Clone Lentiviral Particle - RC212084L2V

Protein Pathways: Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, Jak-STAT signaling

pathway

MW: 57.15 kDa

Gene Summary: The protein encoded by this gene is a cytokine receptor that specifically mediates the

biological effects of interleukin 9 (IL9). The functional IL9 receptor complex requires this protein as well as the interleukin 2 receptor, gamma (IL2RG), a common gamma subunit shared by the receptors of many different cytokines. The ligand binding of this receptor leads to the activation of various JAK kinases and STAT proteins, which connect to different biologic responses. This gene is located at the pseudoautosomal regions of X and Y chromosomes. Genetic studies suggested an association of this gene with the development of asthma. Multiple pseudogenes on chromosome 9, 10, 16, and 18 have been described. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2008]