

Product datasheet for RC212177L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TL1A (TNFSF15) (NM 005118) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: TL1A (TNFSF15) (NM_005118) Human Tagged ORF Clone Lentiviral Particle

Symbol: TL1A

Synonyms: TL1; TL1A; TNLG1B; VEGI; VEGI192A

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 005118

ORF Size: 753 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC212177).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 005118.2

 RefSeq Size:
 2011 bp

 RefSeq ORF:
 756 bp

 Locus ID:
 9966

 UniProt ID:
 095150

 Cytogenetics:
 9q32

 Domains:
 TNF

Protein Families: Druggable Genome, Transmembrane





TL1A (TNFSF15) (NM_005118) Human Tagged ORF Clone Lentiviral Particle - RC212177L1V

Protein Pathways: Cytokine-cytokine receptor interaction

MW: 28.1 kDa

Gene Summary: The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor

(TNF) ligand family. This protein is abundantly expressed in endothelial cells, but is not expressed in either B or T cells. The expression of this protein is inducible by TNF and IL-1 alpha. This cytokine is a ligand for receptor TNFRSF25 and decoy receptor TNFRSF21/DR6. It can activate NF-kappaB and MAP kinases, and acts as an autocrine factor to induce apoptosis in endothelial cells. This cytokine is also found to inhibit endothelial cell proliferation, and thus may function as an angiogenesis inhibitor. Two transcript variants encoding different

isoforms have been found for this gene. [provided by RefSeq, Feb 2011]