

Product datasheet for RC212406L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

WHSC1 (NM_001042424) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: WHSC1 (NM_001042424) Human Tagged ORF Clone Lentiviral Particle

Symbol: WHSC1

Synonyms: KMT3F; KMT3G; MMSET; REIIBP; TRX5; WHS; WHSC1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001042424

ORF Size: 4095 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC212406).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001042424.1</u>

 RefSeq Size:
 7568 bp

 RefSeq ORF:
 4098 bp

 Locus ID:
 7468

 UniProt ID:
 096028

 Cytogenetics:
 4p16.3

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Lysine degradation





ORIGENE

MW: 152.1 kDa

Gene Summary:

This gene encodes a protein that contains four domains present in other developmental proteins: a PWWP domain, an HMG box, a SET domain, and a PHD-type zinc finger. It is expressed ubiquitously in early development. Wolf-Hirschhorn syndrome (WHS) is a malformation syndrome associated with a hemizygous deletion of the distal short arm of chromosome 4. This gene maps to the 165 kb WHS critical region and has also been involved in the chromosomal translocation t(4;14)(p16.3;q32.3) in multiple myelomas. Alternative splicing of this gene results in multiple transcript variants encoding different isoforms. Some transcript variants are nonsense-mediated mRNA (NMD) decay candidates, hence not represented as reference sequences. [provided by RefSeq, Jul 2008]