

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC212523L3V

NFAT4 (NFATC3) (NM_004555) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	NFAT4 (NFATC3) (NM_004555) Human Tagged ORF Clone Lentiviral Particle
Symbol:	NFAT4
Synonyms:	NF-AT4c; NFAT4; NFATX
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_004555
ORF Size:	3204 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC212523).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 004555.2</u>
RefSeq Size:	4005 bp
RefSeq ORF:	3207 bp
Locus ID:	4775
UniProt ID:	<u>Q12968</u>
Cytogenetics:	16q22.1
Domains:	RHD, IPT
Protein Families:	Druggable Genome, Transcription Factors



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US Protein Pathways:Axon guidance, B cell receptor signaling pathway, Natural killer cell mediated cytotoxicity, T
cell receptor signaling pathway, VEGF signaling pathway, Wnt signaling pathwayMW:115.1 kDaGene Summary:The product of this gene is a member of the nuclear factors of activated T cells DNA-binding
transcription complex. This complex consists of at least two components: a preexisting
cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation
and an inducible nuclear component. Other members of this family participate to form this
complex also. The product of this gene plays a role in the regulation of gene expression in T
cells and immature thymocytes. Several transcript variants encoding distinct isoforms have
been identified for this gene. [provided by RefSeq, Nov 2010]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US