

Product datasheet for RC212527L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IGF1 (NM_000618) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: IGF1 (NM_000618) Human Tagged ORF Clone Lentiviral Particle

Symbol: IGF

Synonyms: IGF; IGF-I; IGFI; MGF

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_000618

ORF Size: 459 bp

ORF Nucleotide

TI 005

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC212527).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 000618.2</u>

 RefSeq Size:
 7260 bp

 RefSeq ORF:
 462 bp

 Locus ID:
 3479

 UniProt ID:
 P05019

 Cytogenetics:
 12q23.2

Domains: IIGF

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein





IGF1 (NM_000618) Human Tagged ORF Clone Lentiviral Particle - RC212527L4V

Protein Pathways: Dilated cardiomyopathy, Focal adhesion, Glioma, Hypertrophic cardiomyopathy (HCM), Long-

term depression, Melanoma, mTOR signaling pathway, Oocyte meiosis, p53 signaling pathway, Pathways in cancer, Progesterone-mediated oocyte maturation, Prostate cancer

MW: 17.03 kDa

Gene Summary: The protein encoded by this gene is similar to insulin in function and structure and is a

member of a family of proteins involved in mediating growth and development. The encoded protein is processed from a precursor, bound by a specific receptor, and secreted. Defects in this gene are a cause of insulin-like growth factor I deficiency. Alternative splicing results in multiple transcript variants encoding different isoforms that may undergo similar processing

to generate mature protein. [provided by RefSeq, Sep 2015]