

Product datasheet for RC212571L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL19 (NM_013371) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: IL19 (NM_013371) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL19

Synonyms: IL-10C; MDA1; NG.1; ZMDA1

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_013371

ORF Size: 531 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC212571).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 013371.2

RefSeq Size: 1831 bp
RefSeq ORF: 534 bp
Locus ID: 29949
UniProt ID: Q9UHD0
Cytogenetics: 1q32.1

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway





IL19 (NM_013371) Human Tagged ORF Clone Lentiviral Particle - RC212571L2V

MW: 20.45 kDa

Gene Summary: The protein encoded by this gene is a cytokine that belongs to the IL10 cytokine subfamily.

This cytokine is found to be preferentially expressed in monocytes. It can bind the IL20 receptor complex and lead to the activation of the signal transducer and activator of transcription 3 (STAT3). A similar cytokine in mouse is reported to up-regulate the expression of IL6 and TNF-alpha and induce apoptosis, which suggests a role of this cytokine in inflammatory responses. Alternatively spliced transcript variants encoding the distinct

isoforms have been described. [provided by RefSeq, Jul 2008]