

Product datasheet for RC212619L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ATP2B1 (NM_001682) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: ATP2B1 (NM_001682) Human Tagged ORF Clone Lentiviral Particle

Symbol: ATP2B1

Synonyms: PMCA1; PMCA1kb

Mammalian Cell

Selection:

Puromycin

NM 001682

Vector:

ACCN:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ORF Size: 3660 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC212619).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001682.2

RefSeq Size:6967 bpRefSeq ORF:3663 bpLocus ID:490

 UniProt ID:
 P20020

 Cytogenetics:
 12q21.33

Domains: E1-E2_ATPase, Cation_ATPase_N, Hydrolase, Cation_ATPase_C

Protein Families: Druggable Genome, Transmembrane





ATP2B1 (NM_001682) Human Tagged ORF Clone Lentiviral Particle - RC212619L3V

Protein Pathways: Calcium signaling pathway

MW: 134.7 kDa

Gene Summary: The protein encoded by this gene belongs to the family of P-type primary ion transport

ATPases characterized by the formation of an aspartyl phosphate intermediate during the reaction cycle. These enzymes remove bivalent calcium ions from eukaryotic cells against

very large concentration gradients and play a critical role in intracellular calcium

homeostasis. The mammalian plasma membrane calcium ATPase isoforms are encoded by at

least four separate genes and the diversity of these enzymes is further increased by alternative splicing of transcripts. The expression of different isoforms and splice variants is regulated in a developmental, tissue- and cell type-specific manner, suggesting that these pumps are functionally adapted to the physiological needs of particular cells and tissues. This

gene encodes the plasma membrane calcium ATPase isoform 1. Alternatively spliced

transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul

2008]