

Product datasheet for RC212648L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CAMK2D (NM_001221) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CAMK2D (NM_001221) Human Tagged ORF Clone Lentiviral Particle

Symbol: CAMK2D Synonyms: CAMKD

Mammalian Cell

Puromycin

Selection:

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001221 **ORF Size:** 1497 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC212648).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001221.2</u>

RefSeq Size: 2140 bp
RefSeq ORF: 1500 bp
Locus ID: 817

UniProt ID: Q13557

Cytogenetics: 4q26

Domains: pkinase, TyrKc, S_TKc

Protein Families: Druggable Genome, Protein Kinase





CAMK2D (NM_001221) Human Tagged ORF Clone Lentiviral Particle - RC212648L4V

Protein Pathways: Calcium signaling pathway, ErbB signaling pathway, Glioma, GnRH signaling pathway, Long-

term potentiation, Melanogenesis, Neurotrophin signaling pathway, Olfactory transduction,

Oocyte meiosis, Wnt signaling pathway

MW: 56.2 kDa

Gene Summary: The product of this gene belongs to the serine/threonine protein kinase family and to the

Ca(2+)/calmodulin-dependent protein kinase subfamily. Calcium signaling is crucial for several aspects of plasticity at glutamatergic synapses. In mammalian cells, the enzyme is composed of four different chains: alpha, beta, gamma, and delta. The product of this gene is a delta chain. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Distinct isoforms of this chain have different expression patterns.[provided by

RefSeq, Nov 2008]