

## Product datasheet for RC212775L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## ICAM2 (NM\_001099789) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** ICAM2 (NM\_001099789) Human Tagged ORF Clone Lentiviral Particle

Symbol: ICAM2 Synonyms: CD102

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_001099789

ORF Size: 825 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC212775).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001099789.1

 RefSeq Size:
 1153 bp

 RefSeq ORF:
 828 bp

 Locus ID:
 3384

 UniProt ID:
 P13598

 Cytogenetics:
 17q23.3

**Protein Families:** ES Cell Differentiation/IPS, Transmembrane

**Protein Pathways:** Cell adhesion molecules (CAMs), Natural killer cell mediated cytotoxicity





ORIGENE

**MW:** 30.7 kDa

**Gene Summary:** The protein encoded by this gene is a member of the intercellular adhesion molecule (ICAM)

family. All ICAM proteins are type I transmembrane glycoproteins, contain 2-9 immunoglobulin-like C2-type domains, and bind to the leukocyte adhesion LFA-1 protein. This

protein may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell

adhesion. It mediates adhesive interactions important for antigen-specific immune response,

NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance. Several transcript variants encoding the

same protein have been found for this gene. [provided by RefSeq, Jul 2008]