

## Product datasheet for RC212803L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **DEGS1 (NM\_003676) Human Tagged ORF Clone Lentiviral Particle**

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** DEGS1 (NM\_003676) Human Tagged ORF Clone Lentiviral Particle

Symbol: DEGS1

Synonyms: DEGS; DEGS-1; Des-1; DES1; FADS7; HLD18; MIG15; MLD

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_003676

ORF Size: 969 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC212803).

Sequence:

Cytogenetics:

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 003676.2

 RefSeq Size:
 2101 bp

 RefSeq ORF:
 972 bp

 Locus ID:
 8560

 UniProt ID:
 015121

**Domains:** FA desaturase

**Protein Families:** Druggable Genome, Transmembrane

1q42.11





## DEGS1 (NM\_003676) Human Tagged ORF Clone Lentiviral Particle - RC212803L4V

**Protein Pathways:** Metabolic pathways, Sphingolipid metabolism

**MW:** 37.9 kDa

**Gene Summary:** This gene encodes a member of the membrane fatty acid desaturase family which is

responsible for inserting double bonds into specific positions in fatty acids. This protein contains three His-containing consensus motifs that are characteristic of a group of membrane fatty acid desaturases. It is predicted to be a multiple membrane-spanning protein localized to the endoplasmic reticulum. Overexpression of this gene inhibited biosynthesis of the EGF receptor, suggesting a possible role of a fatty acid desaturase in regulating biosynthetic processing of the EGF receptor. [provided by RefSeq, Mar 2010]