

Product datasheet for RC212807

PKIG (NM 181805) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: PKIG (NM_181805) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: PKIG

Synonyms: PKI-gamma

Mammalian Cell Neomycin

Selection:

Vector:

E. coli Selection: Kanamycin (25 ug/mL)

ORF Nucleotide >RC212807 representing NM_181805

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

pCMV6-Entry (PS100001)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

GATGGGACCACCTCGTCT

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC212807 representing NM_181805

Red=Cloning site Green=Tags(s)

MMEVESSYSDFISCDRTGRRNAVPDIQGDSEAVSVRKLAGDMGELALEGAEGQVEGSAPDKEAGNQPQSS

DGTTSS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6446 g08.zip

Restriction Sites: Sgfl-Mlul



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_181805

ORF Size: 228 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 181805.2</u>

RefSeq Size: 1458 bp
RefSeq ORF: 231 bp
Locus ID: 11142



 UniProt ID:
 Q9Y2B9

 Cytogenetics:
 20q13.12

Protein Families: Druggable Genome


MW: 7.7 kDa

Gene Summary: This gene encodes a member of the protein kinase inhibitor family. Studies of a similar


protein in mice suggest that this protein acts as a potent competitive cAMP-dependent protein kinase inhibitor, and is a predominant form of inhibitor in various tissues. The encoded protein may be involved in osteogenesis. Alternative splicing results in multiple

transcript variants. [provided by RefSeq, Jul 2013]

Product images:



Circular map for RC212807



Western blot validation of overexpression lysate (Cat# [LY416230]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC213682] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).