

Product datasheet for **RC212816L2V**

Activin Receptor Type IIB (ACVR2B) (NM_001106) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Activin Receptor Type IIB (ACVR2B) (NM_001106) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Activin Receptor Type IIB
Synonyms:	ActR-IIB; ACTRIIB; HTX4
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_001106
ORF Size:	1536 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC212816).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_001106.2
RefSeq Size:	1584 bp
RefSeq ORF:	1539 bp
Locus ID:	93
UniProt ID:	Q13705
Cytogenetics:	3p22.2
Domains:	Activin_recp, pkinase, TyrKc, S_TKc



[View online »](#)

Protein Families:	Druggable Genome, Protein Kinase, Transmembrane
Protein Pathways:	Cytokine-cytokine receptor interaction, TGF-beta signaling pathway
MW:	57.5 kDa
Gene Summary:	<p>Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. Type II receptors are considered to be constitutively active kinases. This gene encodes activin A type IIB receptor, which displays a 3- to 4-fold higher affinity for the ligand than activin A type II receptor. [provided by RefSeq, Jul 2008]</p>