

Product datasheet for **RC212849**

Cyclophilin E (PPIE) (NM_203456) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Cyclophilin E (PPIE) (NM_203456) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Cyclophilin E
Synonyms:	CYP-33; CYP33
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC212849 representing NM_203456 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGCCACCACCAAGCGCGTCTTGTACGTGGGTGGACTGGCAGAGGAAGTGGACGACAAAGTTCTTCATG
CTGCGTTCATTCCTTTTGGAGACATCACAGATATTCAGATTCCTCTGGATTATGAAACAGAAAAGCACCG
AGGATTTGCTTTTGTGAATTTGAGTTGGCAGAGGATGCTGCAGCAGCTATCGACAACATGAATGAATCT
GAGCTTTTGGACGTACAATTCGTGTCAATTTGGCCAAACCAATGAGAATTAAGGAAGGCTCTCCAGGC
CAGTTTGGTCAGATGATGACTGGTTGAAGAAGTTTCTGGGAAGACGCTTGAAGAGAATAAAGAGGAAGA
AGGGTCAGAGCCTCCCAAAGCAGAGACCCAGGAGGGAGAGCCCATTTGCTAAAAAGCCCGCTCAAATCCT
CAGGTGTACATGGACATCAAGATTGGGAACAAGCCGGCTGGCCGCATCCAGATGCTCCTGCGTTCTGATG
TCGTGCCCATGACAGCAGAGAATTTCCGCTGCCTGTGCACTCATGAAAAGGGCTTTGGCTTTAAGGGAAG
CAGCTTCCACCGCATCATCCCCAGTTCATGTGCCAGGGCGGTGATTTCAAAACCAATGGCACTGGG
GGCAAGTCCATCTATGGGAAGAAGTTCGATGATGAAAACTTTATCCTCAAGCATAACGGGACCAGGTCTAC
TATCCATGGCCAACTCTGGCCAAACACCAATGGCTCTCAGTTCTTCTGACATGTGACAAGACAGACTG
GCTGGATGGCAAGCATGTGGTGTGGAGAGGTCACCGAAGGCCTAGATGTCTTGGCCAAATTTAGAAA
CAAGAAGAGTCAGCAATTACCAGCCAGCCGAGGTCCTGGAAGCTGACG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC212849 representing NM_203456
Red=Cloning site Green=Tags(s)


MATTKRVLYVGGLAEEVDDKVLHAAFIPFGDITDIQIPLDYETEKHRGF AFVEFELAEDAAAAIDNMNES
 ELFGRTIRVNLAKPMRIKEGSSRPVWSDDDWLKKFSGKLEENKEEEGSEPPKAETQEGEPIAKKARSNP
 QVYMDIKIGNKPAGRIQMLLRSDVVPMTAENFRCLCTHEKGFGFKGSSFHRIIPQFMCQGGDFTNHNGTG
 GKSIYGGKFDDEFILKHTGPGLLSMANSGPNTNGSQFFLTCDKTDWLDGKHVVFGEVTEGLDVLRLQIEK
 QEESAITSQPRSWKLT

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk8050_c04.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_203456

ORF Size: 888 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_203456.2](#), [NP_982281.1](#)

RefSeq Size: 1123 bp

RefSeq ORF: 891 bp

Locus ID: 10450

UniProt ID: [Q9UNP9](#)

Cytogenetics: 1p34.2


Protein Families: Transcription Factors

Protein Pathways: Spliceosome

MW: 32.9 kDa

Gene Summary: The protein encoded by this gene is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein contains a highly conserved cyclophilin (CYP) domain as well as an RNA-binding domain. It was shown to possess PPIase and protein folding activities, and it also exhibits RNA-binding activity. Alternative splicing results in multiple transcript variants. A related pseudogene, which is also located on chromosome 1, has been identified. [provided by RefSeq, Aug 2010]

Product images:



Circular map for RC212849